

REMARK ABOUT THE SPECTRUM OF THE p -FORM LAPLACIAN UNDER A COLLAPSE WITH CURVATURE BOUNDED BELOW

JOHN LOTT

ABSTRACT. We give a lower bound on the number of small positive eigenvalues of the p -form Laplacian in a certain type of collapse with curvature bounded below.

1. INTRODUCTION

A general problem in spectral geometry is to estimate the eigenvalues of the p -form Laplacian on a closed Riemannian manifold M in terms of the geometry of M . From Hodge theory, the number of zero eigenvalues is $b_p(M)$, the p -th Betti number of M . Hence the issue is to understand the positive eigenvalues. The papers [2], [7] and [8] study the case when one assumes an upper bound on the diameter of the manifold and double-sided bounds on the sectional curvatures. An important phenomenon is the possible appearance of positive eigenvalues of the p -form Laplacian that approach zero as a manifold collapses with bounded curvature.

The analysis of [7] and [8] uses the results of Cheeger, Fukaya and Gromov on the geometric structure of manifolds that collapse with double-sided curvature bounds. If one only assumes a lower sectional curvature bound then there are some structure results about collapsing in [4] and [12], but the theory is less developed than in the bounded curvature case.

In this paper we look at the small positive eigenvalues of the p -form Laplacian in an example of collapse with curvature bounded below. Namely, suppose that a compact Lie group G acts isometrically on M on the left. Give G a left-invariant Riemannian metric. For $\epsilon > 0$, let ϵG denote G with its Riemannian metric multiplied by ϵ^2 . Let M_ϵ denote $M = G \backslash (\epsilon G \times M)$ equipped with the quotient Riemannian metric g_ϵ , where G acts diagonally on $\epsilon G \times M$ on the left. If G is connected then $\lim_{\epsilon \rightarrow 0} M_\epsilon = G \backslash M$ in the Gromov-Hausdorff topology, and as ϵ goes to zero, the sectional curvatures of M_ϵ stay uniformly bounded below [12].

For notation, if M is a smooth connected closed manifold with Riemannian metric g , let $\{\lambda_{p,j}(M, g)\}_{j=1}^\infty$ denote the eigenvalues (counted with multiplicity) of the Laplacian on $\overline{\text{Im}}(d) \subset \Omega_{L^2}^p(M)$.

Theorem 1. *If $j = \dim(\text{Ker}(H^p(G \backslash M; \mathbb{R}) \rightarrow H^p(M; \mathbb{R})))$ then $\lim_{\epsilon \rightarrow 0} \lambda_{p,j}(M_\epsilon, g_\epsilon) = 0$.*

In Section 2 we prove Theorem 1. The main points of the proof are the use of a certain variational expression for $\lambda_{p,j}(M, g)$, due to Cheeger and Dodziuk [3], and the avoidance of dealing with the detailed orbit structure of the group action. We then look at the example

Date: February 11, 2002.

1991 *Mathematics Subject Classification.* Primary: 58G25; Secondary: 53C23.

Research supported by NSF grant DMS-0072154.

of an S^1 -action on S^{2n} , which is the suspension of the Hopf action of S^1 on S^{2n-1} , and show that our results slightly improve those of Takahashi [10]. In Section 3 we make some further remarks.

I thank Junya Takahashi for sending me a copy of his paper.

2. PROOF OF THEOREM 1

Let $\mathfrak{g}$ be the Lie algebra of G . It acquires an inner product from the left-invariant Riemannian metric on G . Given $\mathfrak{x} \in \mathfrak{g}$, let $\mathfrak{X}$ be the corresponding vector field on M . Let $i_{\mathfrak{X}}$ denote interior multiplication by $\mathfrak{X}$.

Let $\Omega^*(M)$ denote the smooth differential forms on M . Let $\Omega_{L^2}^*(M)$ be the L^2 -completion of $\Omega^*(M)$. Put

$$\Omega_{max}^*(M) = \{\omega \in \Omega_{L^2}^*(M) : d\omega \in \Omega_{L^2}^{*+1}(M)\}, \quad (2.1)$$

where $d\omega$ is originally defined distributionally.

Put

$$\Omega_G^*(M) = \{\omega \in \Omega^*(M) : g \cdot \omega = \omega \text{ for all } g \in G\} \quad (2.2)$$

and

$$\Omega_{basic}^*(M) = \{\omega \in \Omega_G^*(M) : i_{\mathfrak{X}}\omega = 0 \text{ for all } \mathfrak{x} \in \mathfrak{g}\}. \quad (2.3)$$

Let $\Omega_{G,L^2}^*(M)$ and $\Omega_{basic,L^2}^*(M)$ be the L^2 -completions of $\Omega_G^*(M)$ and $\Omega_{basic}^*(M)$, respectively. Put

$$\Omega_{basic,max}^*(M) = \{\omega \in \Omega_{basic,L^2}^*(M) : d\omega \in \Omega_{basic,L^2}^{*+1}(M)\}, \quad (2.4)$$

where $d\omega$ is originally defined distributionally. Then $\Omega_{basic,max}^*(M)$ is a complex.

From [6] and [11], the cohomology of the complex $\Omega_{basic}^*(M)$ is isomorphic to $H^*(G \backslash M; \mathbb{R})$.

Lemma 1. *The cohomology of the complex $\Omega_{basic,max}^*(M)$ is isomorphic to $H^*(G \backslash M; \mathbb{R})$.*

Proof. The proof is essentially the same as that of [11]. The only point to note is that the homotopy operator A used in the Poincaré lemma in [11] sends $\Omega_{basic,max}^*$ to itself. $\square$

The quotient map $p : \epsilon G \times M \rightarrow M_\epsilon$ defines a principal G -bundle. Pullback gives an isomorphism $p^* : \Omega^*(M_\epsilon) \cong \Omega_{basic}^*(\epsilon G \times M)$. The parallelism of G gives an isomorphism

$$\Omega^*(\epsilon G \times M) \cong C^\infty(G) \otimes \Lambda^*(\mathfrak{g}^*) \otimes \Omega^*(M). \quad (2.5)$$

Taking G -invariants gives isomorphisms

$$\Omega_G^*(\epsilon G \times M) \rightarrow (C^\infty(G) \otimes \Lambda^*(\mathfrak{g}^*) \otimes \Omega^*(M))^G \xrightarrow{\beta} \Lambda^*(\mathfrak{g}^*) \otimes \Omega^*(M), \quad (2.6)$$

where β comes from the map which sends $\sum_k f_k \otimes \eta_k \otimes \omega_k \in C^\infty(G) \otimes \Lambda^*(\mathfrak{g}^*) \otimes \Omega^*(M)$ to $\sum_i f_k(e) \eta_k \otimes \omega_k \in \Lambda^*(\mathfrak{g}^*) \otimes \Omega^*(M)$.

Let $\{\mathfrak{x}_j\}_{j=1}^{dim(G)}$ be a basis of $\mathfrak{g}$. For $\mathfrak{x} \in \mathfrak{g}$, let $e(\mathfrak{x}^*)$ denote exterior multiplication by $\mathfrak{x}^*$ on $\Lambda^*(\mathfrak{g}^*)$.

Lemma 2. *There is an isomorphism of complexes $\mathcal{I} : \Omega^*(M) \rightarrow \Omega_{basic}^*(\epsilon G \times M) \subset \Lambda^*(\mathfrak{g}^*) \otimes \Omega^*(M)$ given by*

$$\begin{aligned} \mathcal{I}(\sigma) &= \left(\prod_{j=1}^{dim(G)} (1 - e(\mathfrak{x}_j^*) \otimes i_{\mathfrak{x}_j}) \right) (1 \otimes \sigma) \\ &= \sum_{k=0}^{dim(G)} (-1)^k \sum_{1 \leq j_1 < \dots < j_k \leq dim(G)} (\mathfrak{x}_{j_k}^* \wedge \dots \wedge \mathfrak{x}_{j_1}^*) \otimes i_{\mathfrak{x}_{j_1}} \dots i_{\mathfrak{x}_{j_k}} \sigma. \end{aligned} \quad (2.7)$$

Proof. If $\sum_k f_k \otimes \eta_k \otimes \omega_k \in (C^\infty(G) \otimes \Lambda^*(\mathfrak{g}^*) \otimes \Omega^*(M))^G$ is G -basic then for $\mathfrak{x} \in \mathfrak{g}$, we also have

$$\sum_k (f_k \otimes i_{\mathfrak{x}} \eta_k \otimes \omega_k + (-1)^{|\eta_k|} f_k \otimes \eta_k \otimes i_{\mathfrak{x}} \omega_k) = 0. \quad (2.8)$$

Then

$$\sum_k (f_k(e) i_{\mathfrak{x}} \eta_k \otimes \omega_k + (-1)^{|\eta_k|} f_k(e) \eta_k \otimes i_{\mathfrak{x}} \omega_k) = 0, \quad (2.9)$$

i.e. if $\sum_k \eta_k \otimes \omega_k$ lies in the image of β restricted to $\Omega_{basic}^*(\epsilon G \times M)$ then

$$\sum_k (i_{\mathfrak{x}} \eta_k \otimes \omega_k + (-1)^{|\eta_k|} \eta_k \otimes i_{\mathfrak{x}} \omega_k) = 0. \quad (2.10)$$

It follows that $\sum_k \eta_k \otimes \omega_k$ can be written as $\mathcal{I}(\sigma)$ for some $\sigma \in \Omega^*(M)$. Thus $\mathcal{I}$ is surjective. It is clearly injective.

It remains to show that $\mathcal{I}$ is a morphism of complexes. Let d^{inv} denote the (finite-dimensional) differential on $\Lambda^*(\mathfrak{g}^*)$. If an element of $\Omega_G^*(\epsilon G \times M)$ is represented as $\sum_k f_k \otimes \eta_k \otimes \omega_k \in C^\infty(G) \otimes \Lambda^*(\mathfrak{g}^*) \otimes \Omega^*(M)$ then the G -invariance implies that for $\mathfrak{x} \in \mathfrak{g}$,

$$\sum_k (\mathfrak{x} f_k \otimes \eta_k \otimes \omega_k + f_k \otimes \eta_k \otimes \mathcal{L}_{\mathfrak{x}} \omega_k) = 0. \quad (2.11)$$

The differential of $\sum_k f_k \otimes \eta_k \otimes \omega_k$ is represented by

$$\sum_k \left(\sum_{j=1}^{dim(G)} \mathfrak{x}_j f_k \otimes e(\mathfrak{x}_j^*) \eta_k \otimes \omega_k + f_k \otimes d^{inv} \eta_k \otimes \omega_k + (-1)^{|\eta_k|} f_k \otimes \eta_k \otimes d\omega_k \right). \quad (2.12)$$

From (2.11), this equals

$$\sum_k \left(- \sum_{j=1}^{dim(G)} f_k \otimes e(\mathfrak{x}_j^*) \eta_k \otimes \mathcal{L}_{\mathfrak{x}_j} \omega_k + f_k \otimes d^{inv} \eta_k \otimes \omega_k + (-1)^{|\eta_k|} f_k \otimes \eta_k \otimes d\omega_k \right). \quad (2.13)$$

Using β , it follows that the induced differential on $\Lambda^*(\mathfrak{g}^*) \otimes \Omega^*(M)$ sends $\sum_k \eta_k \otimes \omega_k$ to

$$\sum_k \left(- \sum_{j=1}^{dim(G)} e(\mathfrak{x}_j^*) \eta_k \otimes \mathcal{L}_{\mathfrak{x}_j} \omega_k + d^{inv} \eta_k \otimes \omega_k + (-1)^{|\eta_k|} \eta_k \otimes d\omega_k \right). \quad (2.14)$$

One can check that when this acts on $\mathcal{I}(\sigma)$, the result is $\mathcal{I}(d\sigma)$. Thus $\mathcal{I}$ is an isomorphism of complexes. $\square$

In fact, under our identifications, $\mathcal{I}$ is the same as p^* .

Let M^{reg} be the union of the principal orbits for the G -action on M . It is a dense open subset of M with full measure. If $m \in M^{reg}$, let $H \subset G$ be its isotropy subgroup, with Lie algebra $\mathfrak{h}$. Define $\alpha : \mathfrak{g} \rightarrow T_m M$ by $\alpha(\mathfrak{x}) = \mathfrak{X}_m$. It passes to an injection $\bar{\alpha} : \mathfrak{g}/\mathfrak{h} \rightarrow T_m M$. For $\epsilon \geq 0$, put $\rho_\epsilon(m) = \det^{1/2}(\epsilon^2 \text{Id}|_{\mathfrak{g}/\mathfrak{h}} + \bar{\alpha}^* \bar{\alpha})$. If $m \notin M^{reg}$, put $\rho_\epsilon(m) = 0$. Note that for $\epsilon > 0$, $\rho_\epsilon^{-1}(m) < \rho_0^{-1}(m)$.

Lemma 3. $\rho_0^{-1} \in L^1(M, d\text{vol})$.

Proof. If $m \in M^{reg}$ then up to an overall constant, $\rho_0(m)$ is the volume of the orbit $G \cdot m$. Then $\int_{M^{reg}} \rho_0^{-1}(m) d\text{vol}(m)$ is proportionate to the volume of $G \backslash M^{reg} \subset G \backslash M$, which is seen to be finite. $\square$

Let $\{\mathfrak{x}_j\}_{j=1}^{\dim(G)}$ be an orthonormal basis of $\mathfrak{g}$.

Lemma 4. For $\epsilon > 0$, there is a positive constant $C(\epsilon)$ such that $\Omega^*(M_\epsilon)$ is isometrically isomorphic to $\Omega^*(M)$ with the new norm

$$\|\omega\|_\epsilon^2 = C(\epsilon) \int_M \rho_\epsilon^{-1}(m) \left(|\omega(m)|_M^2 + \sum_{k=1}^{\dim(G)} \epsilon^{-2k} \sum_{1 \leq j_1 < \dots < j_k \leq \dim(G)} |i_{\mathfrak{x}_{j_1}} \dots i_{\mathfrak{x}_{j_k}} \omega(m)|_M^2 \right) d\text{vol}(m). \quad (2.15)$$

Proof. We can compute the norm squared of $\omega \in \Omega^*(M_\epsilon)$ by taking the local norm squared of $p^*\omega$ on $\epsilon G \times M^{reg}$, dividing by the function which assigns to $(g, m) \in \epsilon G \times M^{reg}$ the volume of the orbit $G \cdot (g, m)$, and integrating over $\epsilon G \times M^{reg}$. If $m \in M^{reg}$ then the relative volume of $G \cdot (g, m)$ is

$$\det^{1/2}(\epsilon^2 \text{Id}|_{\mathfrak{g}} + \alpha^* \alpha) = \epsilon^{\dim(H)} \rho_\epsilon(m). \quad (2.16)$$

The map β of (2.6) is an isometry, up to a constant. As $\{\epsilon^{-1} \mathfrak{x}_j\}_{j=1}^{\dim(G)}$ is an orthonormal basis for $T_\epsilon(\epsilon G)$, the lemma follows from Lemma 2. $\square$

Proof of Theorem 1 : Put $\lambda_{p,j}(\epsilon) = \lambda_{p,j}(M_\epsilon, g_\epsilon)$. From [3],

$$\lambda_{p,j}(\epsilon) = \inf_V \sup_{\eta \in V - 0} \sup_{\theta \in d^{-1}(\eta)} \frac{\|\eta\|_\epsilon^2}{\|\theta\|_\epsilon^2}, \quad (2.17)$$

where V ranges over j -dimensional subspaces of $\text{Im}(d : \Omega^{p-1}(M) \rightarrow \Omega^p(M))$, and $\theta \in d^{-1}(\eta) \subset \Omega^{p-1}(M)$.

Take $j = \dim(\text{Ker}(H^p(G \backslash M; \mathbb{R}) \rightarrow H^p(M; \mathbb{R})))$. From Lemma 1, the inclusion of complexes $\Omega_{basic}^*(M) \rightarrow \Omega_{basic,max}^*(M)$ induces an isomorphism on cohomology. Then there is a j -dimensional subspace V of

$$\text{Ker}(d : \Omega_{basic}^p(M) \rightarrow \Omega_{basic}^{p+1}(M)) \cap \text{Im}(d : \Omega^{p-1}(M) \rightarrow \Omega^p(M)) \quad (2.18)$$

such that if $\eta \in V - 0$ then $\eta \notin \text{Im}(d : \Omega_{basic,max}^{p-1}(M) \rightarrow \Omega_{basic,L^2}^p(M))$. We claim that

$$\lim_{\epsilon \rightarrow 0} \sup_{\eta \in V - 0} \sup_{\theta \in d^{-1}(\eta)} \frac{\|\eta\|_\epsilon^2}{\|\theta\|_\epsilon^2} = 0. \quad (2.19)$$

This will suffice to prove the theorem.

Suppose that (2.19) is not true. Then there are a constant $c > 0$, a sequence $\{\epsilon_r\}_{r=1}^\infty$ in $\mathbb{R}^+$ approaching zero, a sequence $\{\eta_r\}_{r=1}^\infty$ in $V - 0$ and a sequence $\{\theta_r\}_{r=1}^\infty$ in $\Omega^{p-1}(M)$ such that for all r , $d\theta_r = \eta_r$ and

$$\frac{\|\eta_r\|_{\epsilon_r}^2}{\|\theta_r\|_{\epsilon_r}^2} \geq c. \quad (2.20)$$

Doing a Fourier decomposition of θ_r with respect to G , the ratio in (2.20) does not decrease if we replace θ_r by its G -invariant component. Thus we may assume that θ_r is G -invariant.

Without loss of generality, we can replace the norm $\|\cdot\|_\epsilon$ of (2.15) by the same norm divided by $C(\epsilon)$, which we again denote by $\|\cdot\|_\epsilon$. As η_r is smooth on M , it follows from Lemma 3 that the function $\rho_0^{-1}(m) |\eta_r(m)|_M^2$ is integrable on M . Without loss of generality, we may assume that

$$\int_M \rho_0^{-1}(m) |\eta_r(m)|_M^2 d\text{vol}(m) = 1. \quad (2.21)$$

As $\{\eta_r\}_{r=1}^\infty$ lies in the sphere of a finite-dimensional space, there will be a subsequence, which we relabel as $\{\eta_r\}_{r=1}^\infty$, that converges smoothly to some $\eta_\infty \in V - 0$.

From (2.20),

$$\begin{aligned} \|\theta_r\|_{\epsilon_r}^2 &\leq c^{-1} \|\eta_r\|_{\epsilon_r}^2 = c^{-1} \int_M \rho_{\epsilon_r}^{-1}(m) |\eta_r(m)|_M^2 d\text{vol}(m) \\ &\leq c^{-1} \int_M \rho_0^{-1}(m) |\eta_r(m)|_M^2 d\text{vol}(m) = c^{-1}. \end{aligned} \quad (2.22)$$

For large r ,

$$\int_M |\theta_r(m)|_M^2 d\text{vol}(m) \leq (\inf_M \rho_{\epsilon_r}^{-1})^{-1} \int_M \rho_{\epsilon_r}^{-1}(m) |\theta_r(m)|_M^2 d\text{vol}(m) \leq (\inf_M \rho_1^{-1})^{-1} c^{-1}. \quad (2.23)$$

We now work with respect to the metric g on M . By weak-compactness of the unit ball in L^2 , there is a subsequence of $\{\theta_r\}_{r=1}^\infty$, which we relabel as $\{\theta_r\}_{r=1}^\infty$, that converges weakly in L^2 to some $\theta_\infty \in \Omega_{G,L^2}^{p-1}(M)$. Then for $\sigma \in \Omega^p(M)$,

$$\langle \sigma, \eta_\infty \rangle_M - \langle d^* \sigma, \theta_\infty \rangle_M = \lim_{r \rightarrow \infty} (\langle \sigma, \eta_r \rangle_M - \langle d^* \sigma, \theta_r \rangle_M) = \lim_{r \rightarrow \infty} \langle \sigma, \eta_r - d\theta_r \rangle_M = 0. \quad (2.24)$$

Thus $\theta_\infty \in \Omega_{max}^{p-1}(M)$ and $d\theta_\infty = \eta_\infty$.

From (2.22), we also obtain that for each $1 \leq j \leq \dim(G)$,

$$\int_M |i_{\mathfrak{X}_j} \theta_r(m)|_M^2 d\text{vol}(m) \leq (\inf_M \rho_{\epsilon_r}^{-1})^{-1} \int_M \rho_{\epsilon_r}^{-1}(m) |i_{\mathfrak{X}_j} \eta_r(m)|_M^2 d\text{vol}(m) \leq (\inf_M \rho_1^{-1})^{-1} c^{-1} \epsilon_r^2. \quad (2.25)$$

Then for all $\sigma \in \Omega^{p-2}(M)$,

$$\langle \sigma, i_{\mathfrak{X}_j} \theta_\infty \rangle_M = \langle (i_{\mathfrak{X}_j})^* \sigma, \theta_\infty \rangle_M = \lim_{r \rightarrow \infty} \langle (i_{\mathfrak{X}_j})^* \sigma, \theta_r \rangle_M = \lim_{r \rightarrow \infty} \langle \sigma, i_{\mathfrak{X}_j} \theta_r \rangle_M = 0. \quad (2.26)$$

Thus $i_{\mathfrak{X}_j} \theta_\infty = 0$ and $\theta_\infty \in \Omega_{basic,max}^{p-1}(M)$. Hence $\eta_\infty \in \text{Im} \left(d : \Omega_{basic,max}^{p-1}(M) \rightarrow \Omega_{basic,L^2}^p(M) \right)$, which is a contradiction. $\square$

Example : Let $G = U(1)$ act on $M = S^{2n}$ by the suspension of the Hopf action of $U(1)$ on S^{2n-1} . Then $G \backslash M = U(1) \backslash S^{2n}$ is the suspension of $\mathbb{C}P^{n-1}$. One finds that $\text{Ker}(H^p(G \backslash M; \mathbb{R}) \rightarrow H^p(M; \mathbb{R}))$ is nonzero if and only if $p \in \{3, 5, \dots, 2n-1\}$. From Theorem 1, as $\epsilon \rightarrow 0$ there are small eigenvalues of the p -form Laplacian on $\overline{\text{Im}(d)} \subset \Omega_{L^2}^p(M_\epsilon)$ for $p \in \{3, 5, \dots, 2n-1\}$. From the Hodge decomposition, there will also be small eigenvalues of the p -form Laplacian on $\overline{\text{Im}(d^*)} \subset \Omega_{L^2}^p(M_\epsilon)$ for $p \in \{2, 4, \dots, 2n-2\}$. Then using Hodge duality, one concludes that there are small eigenvalues on

1. $\overline{\text{Im}(d^*)} \subset \Omega_{L^2}^1(M_\epsilon)$,
2. $\overline{\text{Im}(d)} \subset \Omega_{L^2}^p(M_\epsilon)$ and $\overline{\text{Im}(d^*)} \subset \Omega_{L^2}^p(M_\epsilon)$ for $p \in \{2, 3, 4, \dots, 2n-3, 2n-2\}$, and
3. $\overline{\text{Im}(d)} \subset \Omega_{L^2}^{2n-1}(M_\epsilon)$.

This slightly sharpens [10, Theorem 1.2]. Note that from eigenvalue estimates for the scalar Laplacian [1], there are no small eigenvalues on $\overline{\text{Im}(d^*)} \subset \Omega_{L^2}^0(M_\epsilon)$, $\overline{\text{Im}(d)} \subset \Omega_{L^2}^1(M_\epsilon)$, $\overline{\text{Im}(d^*)} \subset \Omega_{L^2}^{2n-1}(M_\epsilon)$ or $\overline{\text{Im}(d)} \subset \Omega_{L^2}^{2n}(M_\epsilon)$.

3. REMARKS

1. In the case of a locally-free torus action, there is some intersection between Theorem 1 and the results of [2], [7] and [8]. In [8] one deals with the cohomology of a certain $\mathbb{Z}$ -graded sheaf $H^*(A'_{[0]})$ on the limit space X . In the case of a collapsing coming from a locally-free torus action, Theorem 1 is a statement about the case $* = 0$, when the sheaf $H^0(A'_{[0]})$ is the constant $\mathbb{R}$ -sheaf on X . Of course, the result of Theorem 1 will generally not give all of the small positive eigenvalues that arise in a collapse. As seen in the Example, one can obtain more small eigenvalues just from Hodge duality.

2. Theorem 1 indicates that the relevant cohomology of the limit space is the ordinary cohomology, as opposed for example to the L^2 -cohomology. This is consistent with the results of [8] in the bounded curvature case.

3. If G has positive dimension and acts effectively on M then Theorem 1 describes small positive eigenvalues in a collapsing situation. In some noncollapsing situations, one can show that small eigenvalues do not exist. Here is one such criterion.

Proposition 1. *Let $\mathcal{M}$ be a collection of closed n -dimensional Riemannian manifolds, with $n > 0$. Give $\mathcal{M}$ the Lipschitz metric, coming from biLipschitz homeomorphisms. Suppose that $\mathcal{M}$ can be covered by a finite number of metric balls. For $p \in \mathbb{Z} \cap [0, n]$ and $j \geq 0$, there are positive numbers $a_{p,j}$ and $A_{p,j}$ so that if $(M, g) \in \mathcal{M}$ then $a_{p,j} \leq \lambda_{p,j}(M, g) \leq A_{p,j}$, and $\lim_{j \rightarrow \infty} a_{p,j} = \infty$.*

Proof. Suppose first that for some p and j , there is no upper bound on $\lambda_{p,j}(M, g)$ as (M, g) ranges over $\mathcal{M}$. Then there is a sequence $\{(M_i, g_i)\}_{i=1}^\infty$ in $\mathcal{M}$ with the property that $\lim_{i \rightarrow \infty} \lambda_{p,j}(M_i, g_i) = \infty$. A subsequence of $\{(M_i, g_i)\}_{i=1}^\infty$, which we relabel as $\{(M_i, g_i)\}_{i=1}^\infty$, will have finite distance from some $(M_\infty, g_\infty) \in \mathcal{M}$. Then there are a number $\epsilon \geq 0$ and a sequence of biLipschitz homeomorphisms $h_i : M_\infty \rightarrow M_i$ so that for all i ,

$$e^{-\epsilon} g_\infty \leq h_i^* g_i \leq e^\epsilon g_\infty. \quad (3.1)$$

Here $h_i^* g_i$ is a Lipschitz metric on M_∞ . From Hodge theory,

$$\lambda_{p,j}(M_i, g_i) = \inf_V \sup_{\eta \in V-0} \sup_{\theta \in d^{-1}(\eta)} \frac{\|\eta\|_{M_i}^2}{\|\theta\|_{M_i}^2}, \quad (3.2)$$

where V ranges over j -dimensional subspaces of $\text{Im}(d : \Omega_{max}^{p-1}(M_i) \rightarrow \Omega_{L^2}^p(M_i))$, and $\theta \in d^{-1}(\eta) \subset \Omega_{max}^{p-1}(M_i)$. By naturality,

$$\lambda_{p,j}(M_i, g_i) = \inf_V \sup_{\eta \in V-0} \sup_{\theta \in d^{-1}(\eta)} \frac{\|\eta\|_{h_i^* g_i}^2}{\|\theta\|_{h_i^* g_i}^2}, \quad (3.3)$$

where V ranges over j -dimensional subspaces of $\text{Im}(d : \Omega_{max}^{p-1}(M_\infty) \rightarrow \Omega_{L^2}^p(M_\infty))$, and $\theta \in d^{-1}(\eta) \subset \Omega_{max}^{p-1}(M_\infty)$.

As in [3], it follows from (3.1) and (3.3) that there is a positive integer J which only depends on n so that

$$e^{-J\epsilon} \lambda_{p,j}(M_\infty, g_\infty) \leq \lambda_{p,j}(M_i, g_i) \leq e^{J\epsilon} \lambda_{p,j}(M_\infty, g_\infty). \quad (3.4)$$

This contradicts the assumption that $\lim_{i \rightarrow \infty} \lambda_{p,j}(M_i, g_i) = \infty$.

Now suppose that it is not true that there is a uniform lower bound $a_{p,j}$ on $\{\lambda_{p,j}(M, g)\}_{(M,g) \in \mathcal{M}}$ with the property that $\lim_{j \rightarrow \infty} a_{p,j} = \infty$. Then there are a number $C > 0$, a sequence $\{(M_i, g_i)\}_{i=1}^\infty$ in $\mathcal{M}$ and a sequence of integers $\{j_i\}_{i=1}^\infty$ such that $\lim_{i \rightarrow \infty} j_i = \infty$ and for each i , $\lambda_{p,j_i}(M_i, g_i) \leq C$. Take a subsequence $\{(M_i, g_i)\}_{i=1}^\infty$ and an (M_∞, g_∞) as before. Then for each j ,

$$\lambda_{p,j}(M_\infty, g_\infty) \leq \sup_{i \rightarrow \infty} \lambda_{p,j_i}(M_\infty, g_\infty) \leq \sup_{i \rightarrow \infty} e^{J\epsilon} \lambda_{p,j_i}(M_i, g_i) \leq e^{J\epsilon} C. \quad (3.5)$$

This contradicts the discreteness of the spectrum of the p -form Laplacian on (M_∞, g_∞) . $\square$

Proposition 1 shows that in a certain sense, one has uniform eigenvalue bounds in the noncollapsing case. It seems possible that for a given $n \in \mathbb{Z}^+$, $K \in \mathbb{R}$ and $v, D > 0$, the collection $\mathcal{M}$ of connected n -dimensional Riemannian manifolds (M, g) with sectional curvatures greater than K , volume greater than v and diameter less than D satisfies the hypotheses of Proposition 1. It is known that there is a finite number of homeomorphism types in $\mathcal{M}$ [5]. On the other hand, the analogous space of metrics defined with Ricci curvature, instead of sectional curvature, will generally not satisfy the hypotheses of Proposition 1 [9].

REFERENCES

- [1] P. Bérard, "From Vanishing Theorems to Estimating Theorems: the Bochner Technique Revisited", Bull. Amer. Math. Soc. 19, p. 371-406 (1988)
- [2] B. Colbois and G. Courtois, "A Note on the First Nonzero Eigenvalue of the Laplacian Acting on p -Forms", Manuscripta Math. 68, p. 143-160 (1990)
- [3] J. Dodziuk, "Eigenvalues of the Laplacian on Forms", Proc. Amer. Math. Soc. 85, p. 437-443 (1982)
- [4] K. Fukaya and T. Yamaguchi, "The Fundamental Groups of Almost Nonnegatively Curved Manifolds", Ann. of Math. 136, p. 253-333 (1992)
- [5] K. Grove, P. Petersen and J. Wu, "Geometric Finiteness Theorems via Controlled Topology", Invent. Math. 99, p. 205-213 (1990)
- [6] J. Koszul, "Sur Certains Groupes de Transformations de Lie", in Géométrie différentielle, Colloques Internationaux du CNRS, Strasbourg, p. 137-141 (1953)

- [7] J. Lott, “Collapsing and the Differential Form Laplacian : The Case of a Smooth Limit Space”, to appear, Duke Math. J., <http://www.math.lsa.umich.edu/~lott>
- [8] J. Lott, “Collapsing and the Differential Form Laplacian : The Case of a Singular Limit Space”, preprint, <http://www.math.lsa.umich.edu/~lott>
- [9] G. Perelman, “Construction of Manifolds of Positive Ricci Curvature with Big Volume and Large Betti Numbers”, in Comparison Geometry, Math. Sci. Res. Inst. Publ. 30, Cambridge Univ. Press, Cambridge, p. 157-163 (1997)
- [10] J. Takahashi, “Small Eigenvalues on p -Forms for Collapsing of the Even Dimensional Spheres”, preprint (2001)
- [11] A. Verona, “A de Rham Type Theorem for Orbit Spaces”, Proc. Amer. Math. Soc. 104, p. 300-302 (1988)
- [12] T. Yamaguchi, “Collapsing and Pinching Under a Lower Curvature Bound”, Ann. of Math. 133, p. 317-357 (1991)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109-1109, USA
E-mail address: `lott@math.lsa.umich.edu`